

- rosette assay for the detection of T_{μ} and T_{γ} lymphocytes. *J Immunol Methods* 36:197-209, 1980
10. Jansen J, Schuit HRE, Zwet TL, Meijer CJLM, Hijmans W: Hairycell leukemia: a B-lymphocytic disorder. *Br J Haematol* 42:21-33, 1979
 11. Kung PC, Goldstein G, Reinherz EL, Schlossman SF: Monoclonal antibodies defining distinctive human T-cell surface antigens. *Science* 206:347-349, 1979
 12. Reinherz EL, Kung PC, Goldstein G, Schlossman SF: Further characterization of the human inducer T-cell subset defined by a monoclonal antibody. *J Immunol* 123:2894-2896, 1979
 13. Reinherz EL, Kung PC, Goldstein G, Schlossman SF: A monoclonal antibody reactive with the human cytotoxic/suppressor T-cell subset previously defined by a heteroantiserum termed TH2. *J Immunol* 124:1301-1307, 1980
 14. Warnke R, Levy R: Detection of T- and B-cell antigens with hybridoma monoclonal antibodies: a biotin-avidin-horseradish peroxidase method. *J Histochem Cytochem* 28:771-776, 1980
 15. Warnke R, Miller R, Grogan R, Pederson M, Dilley J, Levy R: Immunologic phenotype in 30 patients with diffuse large cell lymphoma. *N Engl J Med* 303:293-300, 1980
 16. Willemze R, de Graaff-Reitsma CB, Cnossen J, Van Vloten WA, Meijer CJLM: Characterization of T-cell subpopulations in skin and peripheral blood of patients with cutaneous T-cell lymphomas and benign inflammatory dermatoses. *J Invest Dermatol* 80:60-66, 1983
 17. Meijer CJLM, Van der Loo EM, Van Vloten WA: Early diagnosis of mycosis fungoides and Sézary's syndrome by morphometric analysis of lymphoid cells in the skin. *Cancer* 45:2864-2871, 1980
 18. Haynes BF, Metzgar RS, Minna JD, Bunn PA: Phenotype characterization of cutaneous T-cell lymphoma. Use of monoclonal antibodies to compare with other malignant T-cells. *N Engl J Med* 304:1319-1323, 1981
 19. Laroche L, Bach JF: T cell imbalance in nonleukemic and leukemic cutaneous lymphoma defined by monoclonal antibodies. *Clin Immunol Immunopathol* 20:278-284, 1981
 20. Kung PC, Berger CL, Goldstein G, LoGerfo P, Edelson RL: Cutaneous T-cell lymphoma: characterization by monoclonal antibodies. *Blood* 57:261-266, 1981
 21. Fulton R, Thivolet J, Garcier F, Gaucherand M: Les sous-populations de lymphocytes T-helper et suppressor étudiées par les anticorps monoclonaux dans diverses dermatoses. *Ann Dermatol Venerol* 108:243-250, 1981
 22. Zachary CB, MacDonald DM: Quantification of circulating T-lymphocyte subpopulations in atopic eczema using monoclonal antibodies and flow cytofluorimetry. *Br J Dermatol* 107 (suppl 22):14, 1982
 23. Moscicki RA, Morison WL, Parrish JA, Bloch KJ, Colvin RB: Reduction of the fraction of circulating helper-inducer T cells identified by monoclonal antibodies in psoriatic patients treated with long-term psoralen/ultraviolet A radiation (PUVA). *J Invest Dermatol* 79:205-208, 1982
 24. De Pietro WP, Berger CL, Harber LC, Edelson RL: Normal numbers of phenotypic helper, suppressor and total T cell populations in psoriasis vulgaris: quantitation by monoclonal antibodies. *J Am Acad Dermatol* 5:304-307, 1981
 25. Gupta S, Safai B, Good RA: Subpopulations of human T lymphocytes. IV. Quantitation and distribution in patients with mycosis fungoides and Sézary syndrome. *Cell Immunol* 39:18-26, 1978
 26. Munson D, Bunn PA, Schechter GP: Fc receptor defined T-lymphocyte subpopulations in patients with cutaneous T-cell lymphoma. *J Invest Dermatol* 79:317-321, 1981
 27. Thiers BH: Controversies in mycosis fungoides. *J Am Acad Dermatol* 7:1-16, 1982

0022-202X/83/8105-0397\$02.00/0

THE JOURNAL OF INVESTIGATIVE DERMATOLOGY, 81:397-402, 1983

Copyright © 1983 by The Williams & Wilkins Co.

Vol. 81, No. 5

Printed in U.S.A.

Subsets of Epidermal Langerhans Cells as Defined by Lectin Binding Profiles*

GEROLD SCHULER, M.D., NIKOLAUS ROMANI, PH.D., JOHANNA LINERT, ETHAN M. SHEVACH, M.D., AND GEORG STINGL, M.D.

Department of Dermatology, University of Innsbruck (GSch, NR, JL), Innsbruck, Austria; NIAID, National Institutes of Health (EMS), Bethesda, Maryland, U.S.A.; Department of Dermatology I, University of Vienna (GSTi), Vienna, Austria

In this study we characterize the cell surface glycoconjugate moieties of strain 2 guinea pig epidermal Langerhans cells (LC) in single cell suspension by using a battery of 17 fluorescent lectins. All LC displayed binding sites for concanavalin A, succinylated concanavalin

A, *Lens culinaris* agglutinin, *Pisum sativum* agglutinin, wheat germ agglutinin, succinylated wheat germ agglutinin, *Griffonia simplicifolia* agglutinin I, *Ricinus communis* agglutinin I, *Phaseolus vulgaris* E agglutinin, and *Phaseolus vulgaris* L agglutinin, but failed to bind So-

Manuscript received March 31, 1983; accepted for publication June 23, 1983

This work was supported by grant No. 3791 from the Fonds zur Förderung der wissenschaftlichen Forschung, Vienna, Austria.

* This work was presented, in part, at the 11th Annual Meeting of the ESDR, 1981, Amsterdam.

Reprint requests to: Gerold Schuler, M.D., Department of Dermatology, University of Innsbruck, Anichstrasse 35, A-6020 Innsbruck, Austria.

Abbreviations:

anti-Ia.2,4: strain 13 anti-strain 2 serum

EC: epidermal cell(s)

FITC: fluorescein-isothiocyanate

FITC-SpA: FITC-labeled staphylococcal protein A

HRP: horseradish peroxidase

Ia: immune response-associated (antigens)

LC: Langerhans cell(s)

Lectins:

Con A: concanavalin A

DBA: *Dolichos biflorus* agglutinin

GS I-B₄: *Griffonia simplicifolia* I-B₄ isolectin

GSL I: *Griffonia simplicifolia* agglutinin I

HPA: *Helix pomatia* agglutinin

LCA: *Lens culinaris* agglutinin

PHA-E: *Phaseolus vulgaris* E agglutinin

PHA-L: *Phaseolus vulgaris* L agglutinin

PNA: peanut agglutinin

PSA: *Pisum sativum* agglutinin

RCA I: *Ricinus communis* agglutinin I

SBA: soybean agglutinin

SJA: *Sophora japonica* agglutinin

succ. Con A: Succinylated Con A

succ. WGA: succinylated WGA

UEA I: *Ulex europaeus* agglutinin I

WGA: wheat germ agglutinin

PBS-azide: phosphate-buffered saline supplemented with 0.02% sodium azide

TRITC: tetramethylrhodamine-isothiocyanate

phora japonica agglutinin (SJA), *Dolichos biflorus* agglutinin (DBA), and *Ulex europaeus* agglutinin I (UEA I). Neuraminidase pretreatment rendered LC reactive for SJA, but not for DBA and UEA I. The binding profiles of certain lectins point to the existence of LC subpopulations in that *Griffonia simplicifolia* I-B₄ isolectin, peanut agglutinin (PNA), *Helix pomatia* agglutinin, and soybean agglutinin bound to only 80% (range 70–90%) of Ia-positive epidermal cells; binding sites for these lectins on primarily unreactive Ia-positive cells were unmasked when epidermal cells were treated with neuraminidase prior to lectin labeling. Ultrastructural PNA labeling studies revealed that the vast majority of Birbeck granule-containing LC displayed PNA binding sites, whereas indeterminate cells were consistently PNA-negative. Identification of carbohydrate configurations expressed on LC surfaces by lectin binding may provide a clue for the elucidation of the mechanisms of established LC functions and possibly the discovery of as yet unknown properties of this cell type.

It has been widely realized that all cells bear complex carbohydrate moieties at their surface attached both to proteins (forming glycoproteins) and to N-glycosphingosines (forming glycosphingolipids) [1]. Since cell surface carbohydrates play an important role in several specialized cellular functions including receptor activity, cell growth, differentiation, and interaction of cells with their environment [2,3], one may assume that the major biologic role for membrane carbohydrates is to serve as recognition molecules [2,3]. Whereas keratinocyte and melanocyte surface glycoconjugates have at least partly been characterized [4–22], no information at all exists on the surface carbohydrate composition of epidermal LC.

Lectins have emerged as a valuable tool for the investigation of the complex carbohydrates associated with cell surfaces [23,24] by their binding to specific sugar residues [25]. In this study we investigated the cell surface glycoconjugate pattern of guinea pig epidermal LC by using a battery of 17 lectins which recognize the majority of sugars usually present in membrane glycoproteins [26] and glycolipids [27].

MATERIALS AND METHODS

Animals

Initial breeding pairs of inbred strain 2 and strain 13 guinea pigs were obtained from breeding colonies of the National Institutes of Health, Bethesda, Maryland.

Antisera

The major histocompatibility complex of strain 2 guinea pigs differs from that of strain 13 guinea pigs only in the I-region [28]. Strain 13 anti-strain 2 (anti-Ia.2,4) serum was prepared as previously described [29]. Before use antiserum was inactivated at 56°C for 30 min.

Preparation of Epidermal Cell Suspensions

Single cell suspensions of strain 2 guinea pig epidermal cells (EC) were prepared from white-spotted abdominal skin by standard trypsinization procedures according to the method of Stingl et al [30].

Fluorescent Lectin Labeling

Reagents: TRITC-labeled lectins were purchased from Vector Laboratories (Burlingame, California), and E-Y Laboratories Inc. (San Mateo, California) and FITC-labeled GS I-B₄, PNA, SBA, and HPA from E-Y Laboratories Inc.; inhibitory sugars were obtained from Sigma Co. (St. Louis, Missouri).

Labeling Techniques:

(a) **Fluorescent lectin staining reactions for visualization of cell surface glycoconjugates:** Lectin labeling was carried out on unfixed cells to rule out possible fixation-induced alterations of cell surface glycoconjugates, in particular the exposure of masked lectin receptors. Artifacts possibly induced by redistribution phenomena and endocytic uptake of the

lectins were prevented by performing incubation and rinsing procedures at 4°C with 0.02% sodium azide added to all solutions. Optimal working concentrations of labeled lectins were determined in pilot experiments as described previously [21]. For lectin labeling, 250 μ l of precooled EC suspensions (resuspended to 4×10^6 EC/ml PBS-azide) were added to equal volumes of conjugated lectins diluted in PBS-azide to give a final concentration of 12.5–100 μ g/ml (depending on the particular lectin); the reaction mixture was incubated at 4°C for 20 min, cells were then pelleted by centrifugation and washed 5 times with cold PBS-azide.

(b) **Indirect immunofluorescence technique for identification of Ia-positive cells (i.e., LC):** Ia-positive EC (i.e., LC) [30] were identified by an indirect immunofluorescence procedure using anti-Ia.2,4 serum and FITC-SpA as described previously [30]. Since GS I-B₄ isolectin, PNA, SBA, and HPA were available only as FITC conjugates, TRITC-labeled rabbit F(ab')₂ anti-guinea pig IgG diluted 1:20 with PBS-azide (Cappel Laboratories, Cochranville, Pennsylvania) was used instead of FITC-SpA for identification of Ia-positive cells.

Labeling Sequence: With each experiment separate samples of EC suspensions were subjected to: (1) fluorescent lectin labeling (= procedure a) followed by staining for Ia antigens (= procedure b) and to (2) Ia staining followed by lectin labeling. The strong reactivity of certain lectins (Con A, WGA, SBA, GSL I, RCA I, GS-I B₄) led frequently to agglutination of resuspended EC. To enable unequivocal evaluation of LC labeling profiles, EC were first attached (20 min, 4°C, PBS-azide) to poly-L-lysine-coated glass slides [31] and then labeled as described above. EC were used unfixed as well as prefixed (1% paraformaldehyde in PBS for 30 min at 4°C followed by exposure to 10 mM NH₄Cl in PBS for quenching of unreacted aldehyde groups) to check for possible redistribution phenomena of surface molecules following attachment.

Neuraminidase Pretreatment

For nonreactive lectins the binding capacity of LC was investigated after neuraminidase pretreatment. Unfixed EC were exposed for 60 min at 37°C to *Vibrio comma* neuraminidase (Behringwerke, Marburg, F.R.G.) at a concentration of 0.01 IU/ml in serum-free Medium 199, thoroughly rinsed with PBS, and then subjected to the labeling procedures as outlined above.

Control Experiments

Controls carried along with each experiment included: (1) incubation of cells with lectin conjugates in the presence of 0.1 M inhibitory sugar (see Table I) for determination of lectin binding specificity. In the case of PHA-E and PHA-L reacting with complex carbohydrate configurations [25], appropriate haptenic oligosaccharides were not available. (2) to rule out false negative lectin binding resulting from possible nonspecific action of the sugar on cell surface moieties (e.g., shedding of glycoconjugates), cells were also incubated in the presence of non-inhibitory sugar at 0.1 M concentration. (3) substitution of anti-Ia.2,4 serum by normal guinea pig serum as a negative control for the indirect immunofluorescence procedure detecting Ia-positive cells. (4) incubation of EC with TRITC-labeled lectins followed by FITC-SpA to exclude a possible affinity of FITC-SpA for lectin-treated cells; FITC-GS I-B₄ stained EC were incubated with TRITC rabbit F(ab')₂ anti-guinea pig IgG.

Ultrastructural Lectin Staining

Reagent: HRP-conjugated PNA was purchased from E-Y Laboratories Inc.

Labeling procedure: Determination of the optimal lectin concentration was performed as described in detail elsewhere [21]. Two hundred microliters of EC suspension (4×10^6 /ml PBS) were added to equal volumes of HRP-conjugated PNA to give a final concentration of 200 μ g/ml. The controls were incubated with the respective lectin in the presence of the specific inhibitory sugar (0.1 M). After incubating for 60 min on ice, cells were pelleted and washed 3 times with ice-cold PBS, fixed with precooled half-strength Karnovsky's fixative [32] for 60 min at 20°C, rinsed with 0.1 M cacodylate buffer (pH 7.4) followed by 0.05 M Tris-HCl buffer (pH 7.6), and subjected to the DAB reaction (20 min, 20°C in the dark) [33]. After rinsing procedures, cells were filtered onto Unipore polycarbonate membranes (pore size 0.2 μ m, Biorad Laboratories, Richmond, California); to avoid cell losses the filters were then carefully covered with a 2% agarose solution, fixed for another 30 min in half-strength Karnovsky's fixative, and processed according to our standard embedding procedures [21]. Ultrathin sections were cut through the depth of the pellet, collected on hexagonal grids, and examined with a Philips EM 400 electron microscope operating at 80 kV.

Ultrastructural Identification of EC

Keratinocytes (K) were identified by bundles of tonofilaments and melanosome complexes. LC were identified by the established morphologic criteria, in particular by the presence of Birbeck granules [34]. Cells apparently lacking these organelles even on several consecutive sections but displaying all other morphologic criteria of LC were rated as "indeterminate" cells. Since these cells are known to express identical surface markers as Birbeck granule-containing LC, they are considered to belong to the LC lineage [35]. In order to ascertain that these indeterminate cells bear Ia antigens, a portion of the EC samples was processed for the immunoelectron microscopic demonstration of Ia antigens. EC suspensions (20×10^6 /ml PBS-azide) were incubated for 30 min at 4°C with an equal amount of anti-Ia-2.4 serum (diluted 1:5), or with heat-inactivated serum from nonimmunized strain 13 guinea pigs for control purposes, washed 3 times, resuspended (5×10^6 /ml PBS-azide), and reacted with an equal amount of protein A-iron-dextran (1 mg/ml) [36] for 30 min on ice, fixed in half-strength Karnovsky's fixative (60 min, 20°C) [32], and processed according to our routine embedding procedures [21].

RESULTS

General

Identical lectin binding patterns of LC were observed when lectin staining was either preceded or followed by Ia staining (see *Materials and Methods*). Similarly, attached EC and EC in suspension displayed identical lectin binding profiles, which were also not altered by prefixation of EC or by the presence of noncompeting sugars. All the other controls (see *Materials and Methods*) were consistently negative.

Lectin Binding Profiles of LC

According to their different binding patterns to LC, the lectins employed in this study can be divided into the following three categories (Table I):

Group 1: Lectins binding to all LC—Ten of the lectins reacted with all Ia-positive cells (i.e., LC) in a given suspension (Table I). These lectins reacted also with the majority of other EC but with varying degrees of intensity.

Group 2: Lectins unreactive with LC—Three of the lectins used conformed to this binding pattern (Table I). Whereas

UEA I did not bind to any EC even after neuraminidase pretreatment (but reacted strongly with human O erythrocytes used as a positive control), DBA and SJA reacted with a considerable portion of EC, but did not bind to any LC. Neuraminidase pretreatment was performed in order to determine whether this lack of reactivity was due to either an absence or an inaccessibility of the respective sugar residues. Such a treatment failed to uncover DBA binding sites, but rendered LC reactive for SJA (Table I). These results indicate a true absence of DBA binding sites and suggest that SJA receptors are naturally masked by sialic acid on LC.

Group 3: Lectins binding to subfractions of LC—Four lectins conformed to this binding pattern in that GS I-B₄, PNA, SBA, and HPA bound only to approximately 80% (range 70–90%) of Ia-positive EC (Table I). LC reactive with PNA, SBA, and HPA displayed uniform fluorescence intensity, whereas among GS I-B₄ positive LC, we observed an inverse relationship between cell size and lectin staining intensity (Table I). Ia-positive EC unreactive with either GS I-B₄, PNA, SBA, or HPA in general were smaller than those possessing binding sites for these lectins (Figs 1, 2). Neuraminidase pretreatment induced binding sites for all group 3 lectins on initially unreactive LC; unmasking of the respective receptors was apparent with the lectins SBA, HPA, and PNA, but less obvious with GS I-B₄.

Ultrastructural PNA labeling revealed important differences between LC and indeterminate cells. Whereas the majority of Birbeck granule-containing LC displayed PNA binding sites, indeterminate cells (i.e., cells exhibiting all morphologic criteria of LC but apparently devoid of Birbeck granules) were consistently unreactive for PNA (Fig 3). Both cell populations, however, were uniformly Ia-positive as demonstrated by immunoelectron microscopy (not shown).

DISCUSSION

LC are dendritic epidermal cells and play a functional role in the afferent limb of the immune response which is similar to that of Ia-bearing macrophages [37]. Since it is very likely that—analogueous to macrophages [38]—many of the functional capacities of this important cell type [39] depend on cell surface

TABLE I. Lectin labeling profiles

Lectins	Percentage of lectin binding EC		Percentage of lectin binding LC	
	Untreated	Neuraminidase	Untreated	Neuraminidase
Group 1				
Con A	100 (++++) ^a	n.d. ^b	100 (+++)	n.d.
succ. Con A	100 (+)	n.d.	100 (+)	n.d.
LCA	95 (+/+++)	n.d.	100 (++)	n.d.
PSA	95 (+/+++)	n.d.	100 (++)	n.d.
WGA	99 (++)/++++)	n.d.	100 (++)	n.d.
succ. WGA	78 (+)	n.d.	100 (+)	n.d.
GSL I	95 (++)/++++)	n.d.	100 (++)	n.d.
RCA I	99 (++)/++++)	n.d.	100 (++)	n.d.
PHA-E	100 (++)/++++)	n.d.	100 (++)	n.d.
PHA-L	100 (++)/++++)	n.d.	100 (++)	n.d.
Group 2				
SJA	55 (+)	100 (++)/++++)	0	100 (++)
DBA	25 (+)	40 (++)	0	0
UEA I	0	0	0	0
Group 3				
GSI-B ₄	90 (+/+++)	100 (++)/++++)	75–90 (+/+++)	100 (++)/++++)
PNA	94 (+/+++)	100 (++)/++++)	70–90 (+)	100 (++)
HPA	25 (+)	100 (++)	70–90 (+)	100 (++)
SBA	85 (+/+++)	97 (++)/++++)	65–85 (+)	100 (++)

^a Fluorescence intensity: expressed semiquantitatively using a scale range from – to +++.

^b n.d. = not determined.

Major sugar specificities [25]: Con A, succ. Con A, LCA, PSA: α -D-mannosyl, α -D-glucosyl; WGA: (β -N-acetylglucosaminyl)-n, sialic acid; succ. WGA: β -N-acetylglucosaminyl-n; GSL I: α -D-galactosyl and N-acetyl- α -D galactosaminyl end groups; RCA I: β -D-galactosyl; PHA-E, PHA-L: complex carbohydrate binding specificity; SJA: N-acetyl- β -D-galactosaminyl; SBA, DBA, HPA: N-acetyl- α -D-galactosaminyl; UEA I: α -L-fucosyl; GSI-B₄: α -D-galactosyl end groups; PNA: β -D-galactosyl-(terminal).

Inhibitory sugars (0.1 M): Con A, succ. Con A: α -methyl-D-glucopyranoside; LCA, PSA: α -methyl-D-mannopyranoside; WGA, succ. WGA: N'-N'-diacetyl chitobiose; GSL I: α -D-meliobiose; RCA I: lactose; PHA-E, PHA-L: no inhibitory sugar available; SJA, DBA, HPA, SBA: N-acetyl-D-galactopyranoside; UEA I: L-fucose; GSI-B₄, PNA: α -methyl-D-galactopyranoside.




FIG 1 and 2. EC suspensions were labeled with FITC-conjugated GS I-B₄ lectin followed by Ia staining (indirect immunofluorescence technique employing TRITC-labeled second antibody) for identification of LC. A particular field was photographed by plain light microscopy (B), FITC (A), and TRITC (C) filter setting. $\times 800$. Fig 1: Arrows denote LC (C) only slightly reactive with FITC GS I-B₄ (A). Fig 2: Arrows denote a small and a medium sized LC unreactive and strongly reactive for FITC GS I-B₄ lectin (A), respectively.

interactions, we have been interested in defining the complex surface carbohydrate moieties by using lectins as specific probes. We elected to study EC suspensions rather than frozen skin sections for several reasons: first, identification of LC and their surface glycoconjugates is easier; secondly, interpretation is more accurate due to the absence of the constituents of the extracellular space; and thirdly, all in vitro studies on the functional capabilities of LC have also employed this model system.

The binding of group 1 lectins to LC surfaces is not surprising since the respective carbohydrate residues commonly are found on most cell types so far investigated, including keratinocytes and melanocytes [1,4–22].

Neuraminidase pretreatment of EC showed that the failure of certain lectins (group 2) to bind to Ia-positive EC may have different reasons. Since neuraminidase failed to uncover UEA I and DBA receptors on LC, one may suggest that the respective binding sites are either absent or present only in such minute amounts that they escape detection by the methods used; alternatively, it is conceivable that these residues are sterically inaccessible due to glycosylation of neighboring sugar residues [25]. In contrast, neuraminidase pretreatment of EC enables LC to bind SJA, which indicates that the respective receptor sites are present, but normally masked by sialic acid—a phenomenon by no means unique to LC [21,22,40–44].

By far the most interesting finding arising from the present work is the existence of lectins (group 3) which react only with certain LC subsets. The additional finding that neuraminidase pretreatment restored the capacity of LC to bind all group 3 lectins suggests that cell membranes of LC differ in their degree of sialylation.

Though we got the impression that LC unreactive with GS I-B₄, PNA, SBA, and HPA were not only smaller but generally also less Ia-positive than LC reactive with these lectins, flow cytofluorometric studies are required to confirm these observations. Double-labeling experiments at the ultrastructural level are needed to determine to what extent group 3 lectins identify overlapping LC populations.

A striking observation was that indeterminate cells consistently lacked PNA binding sites whereas the majority of Birbeck granule-containing LC proved to be PNA reactive. Whether these differences in lectin binding profiles reflect the existence of functionally diverse LC subsets (including indeterminate cells) is as yet unknown but appears conceivable in that differentiation and maturation [42,45,46], activation [47], homing mechanisms [48], receptor function [49], and immune recognition [50] of immunoactive cells are critically linked to the quantity and quality of sugar moieties expressed on their cell membrane. Although according to current understanding, cells of the LC lineage are the only Ia-positive cells within normal guinea pig epidermis [30,35,39], one should even consider the theoretical possibility that a minor portion of the Ia-positive EC (e.g., those that fail to bind group 3 lectins) might represent a hitherto unrecognized, functionally distinct EC population. In order to determine whether subfractions of Ia-positive EC as defined by different lectin binding patterns differ in their capacity to induce antigen-specific and/or allogeneic T-cell activation [37], we are currently using group 3 lectins for the fractionation [51] of highly enriched LC suspensions [52].

In summary, identification of carbohydrate configurations expressed on LC surfaces by lectin binding has revealed the existence of LC subpopulations and may provide a clue for the




FIG 3. Ultrastructural PNA labeling. *A* and *C*, An epidermal Langerhans cell (*LC*) identified by the presence of Birbeck granules (*C*, arrows), as well as an adjacent keratinocyte (*t* = tonofilaments) display PNA binding sites on their cell membranes as evidenced by deposition of HRP reaction product (*A*, *C*). *B* and *D*, No PNA reactivity is disclosed on an indeterminate cell (*IC*) (*B*, *D*), whereas an adjacent keratinocyte possesses PNA binding sites as indicated by deposition of HRP reaction product. Bars = 2 μ (*A*, *B*) and 0.5 μ (*C*, *D*).

elucidation of the mechanisms of established LC functions and possibly the discovery of as yet unknown properties of this cell type.

We are grateful to Mrs. Janet Gschnitzer for excellent secretarial assistance and to Mr. Bernhard Sickert for his expert photographic work. We thank Dr. R.S. Molday, Vancouver for his generous gift of protein A-iron-dextran.

REFERENCES

1. Gahmberg CG: Membrane glycoproteins and glycolipids: structure, localization and function of the carbohydrate, *New Comprehensive Biochemistry* (general editors: A Neuberger, LLM Van Deenen), vol 1, Membrane Structure. Edited by JB Finean, RH Michell. Amsterdam/New York/Oxford, Elsevier/North-Holland, 1981, pp 127-160
2. Olden K, Parent JB, White SL: Carbohydrate moieties of glycoproteins. A re-evaluation of their function. *Biochim Biophys Acta* 650:209-232, 1982.
3. Hakomori S: Glycosphingolipids in cellular interaction, differentiation, and oncogenesis. *Annu Rev Biochem* 50:733-764, 1981
4. Nieland ML: Epidermal intercellular staining with fluorescein-conjugated phytohemagglutinins. *J Invest Dermatol* 60:61-66, 1973
5. Hashimoto K, King LE, Yamanishi Y, Beachey EH, Maeyens EJ: Identification of the substance binding pemphigus antibody and

- concanavalin A in the skin. *J Invest Dermatol* 62:423-435, 1974
6. Van Lis MJM, Kalsbeek GL: The interaction of concanavalin A and the surface coat of stratified squamous epithelium. *Br J Dermatol* 92:27-35, 1975
7. Holt PJA, Anglin JH, Nordquist JR, Nordquist RE: Localization of specific carbohydrate configurations in human skin using fluorescein-labeled lectins. *Br J Dermatol* 100:237-245, 1979
8. Nemanic MK, Elias PM: Localization and identification of sugars in mammalian epidermis. *J Cell Biol* 83:46a, 1979
9. Galoppin L, Saurat JH: Dynamic redistribution of concanavalin A binding sites on isolated guinea pig keratinocytes. *Arch Dermatol Res* 265:63-69, 1979
10. Brabec RK, Peters BP, Bernstein IA, Gray RH, Goldstein IJ: Differential lectin binding to cellular membranes in the epidermis of the newborn rat. *Proc Natl Acad Sci USA* 77:477-479, 1980
11. King IA, Tabiowo A, Williams RH: Incorporation of L-(³H) fucose and D-(³H) glucosamine into cell surface associated glycoconjugates in epidermis of cultured pig skin slices. *Biochem J* 190:65-77, 1980
12. Mann PR, Williams RH, Gray GM: Distribution of glycoprotein containing fucose in normal and psoriatic keratinocytes. *Br J Dermatol* 102:649-657, 1980
13. Gommans JM, Van den Hurk JJMA: Lectin binding studies on suspensions of isolated epidermal keratinocytes. *Br J Dermatol* 104:641-648, 1981
14. Davies HW, Trotter MD: Synthesis and turnover of membrane glycoconjugates in monolayer culture of pig and human epidermal cells. *J Invest Dermatol* 104:649-658, 1981
15. Reano A, Faure M, Jacques Y, Reichert U: Use of lectins for the study of human epidermal differentiation. *Br J Dermatol* 107(suppl 23):143-146, 1982
16. Brysk MM, Snider JM: Concanavalin A binding glycoproteins of epidermal cells. *J Invest Dermatol* 79:193-197, 1982
17. Brysk MM, Snider JM: The effect of the state of differentiation on labeling of epidermal cell surface glycoproteins. *J Invest Dermatol* 78:366-370, 1982
18. Reano A, Faure M, Jacques Y, Reichert U, Schaefer H, Thivolet J: Lectins as markers of human epidermal cell differentiation. *Differentiation* 22:205-210, 1982
19. Zieske JK, Bernstein IA: Modification of cell surface glycoprotein: addition of fucosyl residues during epidermal differentiation. *J Cell Biol* 95:626-631, 1982
20. Rittmann BR, Mackenzie IC, Rittman GA: Lectin binding to murine oral mucosa and skin. *Arch Oral Biol* 27:1013-1021, 1982
21. Schuler G, Romani N, Fritsch PO: Differential lectin binding to mammalian keratinocytes and melanocytes in vitro. *Eur J Cell Biol* 27:88-95, 1982
22. Romani N, Schuler G, Fritsch P: Identical lectin binding patterns of human melanocytes and melanoma cells in vitro. *J Invest Dermatol* 80:272-277, 1983
23. Sharon N, Halina HS: Use of lectins for the study of membranes, *Methods in Membrane Biology*, vol 3. Edited by ED Korn. New York, Plenum Press, 1975, pp 147-200
24. Lotan R: Qualitative and quantitative aspects of labeling cell surface carbohydrates using lectins as probes, *Cell Surface Labeling*. Edited by RP Becker, O Johari. Scanning Electron Microscopy Inc., 1979, pp 549-564
25. Goldstein IJ, Hayes CE: The lectins: carbohydrate binding proteins of plants and animals. *Adv Carb Chem Biochem* 35:127-340, 1978
26. Kornfeld R, Kornfeld S: Comparative aspects of glycoprotein structure. *Annu Rev Biochem* 45:217-237, 1976
27. Laine RA, Stellner K, Hakomori S: Isolation and characterization of membrane glycosphingolipids. *Methods Membrane Biol* 2:205-243, 1974
28. Geczy AF, de Weck AL, Schwartz BD, Shevach EM: The major histocompatibility complex of the guinea pig. I. Serologic and genetic studies. *J Immunol* 115:1704-1710, 1975
29. Shevach EM, Rosenstreich DL, Green I: The distribution of histocompatibility antigens on T and B cells in the guinea pig. *Transplantation* 16:126-133, 1973
30. Stingl G, Katz SI, Shevach EM, Wolff-Schreiner E, Green I: Detection of Ia antigens on Langerhans cells in guinea pig skin. *J Immunol* 120:570-578, 1978
31. Mazia D, Schatten G, Sale W: Adhesion of cells to surfaces coated with poly-lysine: applications to electron microscopy. *J Cell Biol* 66:198-200, 1975
32. Karnovsky MJ: A formaldehyde-glutaraldehyde fixative of high osmolality for use in electron microscopy. *J Cell Biol* 27:137A-138A, 1965
33. Graham RC Jr, Karnovsky MJ: The early stages of absorption of injected horseradish peroxidase in the proximal tubules of mouse kidney. Ultrastructural cytochemistry by a new technique. *J Histochem Cytochem* 14:291-302, 1966
34. Wolff K: The Langerhans cell. *Curr Probl Dermatol* 4:79-145, 1972
35. Stingl G: New aspects of Langerhans cell function. *Int J Dermatol* 19:189-213, 1980
36. Molday RS, Mackenzie D: Immunospecific ferromagnetic iron-dextran reagent for the labeling and magnetic separation of cells. *J Immunol Methods* 52:353-367, 1982
37. Stingl G, Katz SI, Shevach EM, Rosenthal AS, Green I: Analogous functions of macrophages and Langerhans cells in the initiation of the immune response. *J Invest Dermatol* 71:59-64, 1978
38. Rosenstreich DL: *The Macrophage, Cellular Functions in Immunity and Inflammation*. Edited by JJ Oppenheim, DL Rosenstreich, M Potter. New York/Amsterdam, Elsevier North Holland, 1981, pp 127-159
39. Stingl G, Wolff K: *The Langerhans cell, Update, Dermatology in General Medicine*. Edited by TB Fitzpatrick, AZ Eisen, K Wolff, IM Freedberg, KF Austen. New York, McGraw-Hill 1983, pp 1-12
40. Goldman R, Sharon N, Lotan R: A differential response elicited in macrophages on interaction with lectins. *Exp Cell Res* 99:408-422, 1976
41. Pfenninger KH, Maylié-Pfenninger MF: Lectin labeling of sprouting neurons I. Regional distribution of surface glycoconjugates. *J Cell Biol* 89:536-546, 1981
42. Sharon N: Cell surface receptors for lectins: markers of murine and human lymphocyte subpopulations, 4th International Congress of Immunology 80, *Progress in Immunology IV*, vol 1. Edited by M Fougereau, J Dausset. London, Academic Press, 1980, pp 254-273
43. Skutelsky E, Lotan R, Sharon N, Danon D: Distribution of the T-antigen on erythroid cell surfaces. Studies with peanut agglutinin, an anti-T specific lectin. *Biochim Biophys Acta* 467:165-174, 1977
44. Maylié-Pfenninger MF, Jamieson JD: Distribution of cell surface saccharides on pancreatic cells. II. Lectin labeling patterns on mature guinea pig and rat pancreatic cells. *J Cell Biol* 80:77-95, 1979
45. Feizi T: Carbohydrate differentiation antigens. *Trends in Biochemical Sciences* 6:333-335, 1981
46. Haimovitz A, Fuks Z, Galili N, Treves AJ: Changes in peanut agglutinin binding to human monocytes during their maturation to macrophages. *J Reticuloendothelial Soc* 31:187-192, 1982
47. Maddox DE, Shibita S, Goldstein IJ: Stimulated macrophages express a new glycoprotein receptor reactive with *Griffonia simplicifolia* I-B₄ isolectin. *Proc Natl Acad Sci USA* 79:166-170, 1982
48. Woodruff JJ, Gesner B: The effect of neuraminidase on the fate of transfused lymphocytes. *J Exp Med* 129:551-567, 1969
49. Poste G, Allen H, Matta KL: Cell surface receptors for lymphokines. II. Studies on the carbohydrate composition of the MIF receptor on macrophages using synthetic saccharides and plant lectins. *Cell Immunol* 44:89-98, 1979
50. Mann DL, Muchmore A: Inhibition of DRw antisera reactivity by mono- and oligosaccharides. *J Immunol* 124:2218-2221, 1980
51. Reisner Y, Sharon N: Cell fractionation by lectins. *Trends in Biochemical Sciences* 5:29-31, 1980
52. Schuler G, Auböck J, Linert J: Enrichment of epidermal Langerhans cells by immunoadsorption to *Staphylococcus aureus* cells. *J Immunol* 130:2008-2010, 1983